

**TABLE 1.** Clinicopathologic characteristics of 43 patients who underwent pancreatectomy with PV/SMV resection for BR/LA PC.

		n (%)
Gender	Male	24 (55.8)
	Female	19 (44.2)
Age, years [median (IQR)]		65 (57-70)
ECOG	0	19 (44.2)
	1	14 (32.5)
	2	10 (23.3)
Location	Body	10 (23.3)
	Head	8 (18.6)
	Neck	12 (27.9)
	Uncinate	13 (30.2)
Neoadjuvant Chemotherapy	No	17 (39.5)
	Yes	26 (60.4)
Operation	Total Pancreatectomy	23 (53.4)
	Whipple	13 (30.2)
	Distal Pancreatectomy	7 (16.2)
Venous Reconstruction Type	Primary	17 (39.5)
	PTFE	18 (41.8)
	Lateral Venorhography	8 (18.6)
Length resected, cm [median (IQR)]		3 (2-4)
ICU stay	No	33 (76.7)
	Yes	10 (23.2)
ICU stay, days [median (IQR)]		0 (0-0)
Transfused pRBC units [median (IQR)]		2 (0-3)
Postoperative hospital LoS, days [median (IQR)]		9 (7-14)
Adjuvant Chemotherapy	No	27 (64.2)
	Yes	15 (35.7)
Tumour size, cm [median (IQR)]		3.2 (2.5-4.2)
T	T1	8 (18.6)
	T2	15 (34.8)
	T3	18 (41.8)
	T4	2 (4.6)
N	N0	7 (16.2)
	N1	30 (69.7)
	N2	6 (14)
Resection	R0	37 (86)
	R1	6 (14)
	R2	0 (0)
Total LNs [median (IQR)]		27 (18-35)
Positive LNs [median (IQR)]		2 (1-5)
LN ratio, % [median (IQR)]		8 (4-20)
Vein Infiltration	No	19 (44.2)
	Yes	24 (55.8)
Complications	No	31 (72)
	Yes	12 (28)
Reoperation	No	38 (88.3)
	Yes	5 (11.4)

ICU: intensive care unit, LOS: length of stay, LN: lymph node.